

Medical Herbalism

A Journal for the Clinical Practitioner

Volume 13, Number 2

Winter 2002-2003

Herbs and insulin resistance

by Paul Bergner

Abstract: The increasing recognition of the pathological effects of hyperinsulinemia with and without loss of glycemic control has led to the search for agents that increase insulin sensitivity in preference to those that stimulate insulin secretion or inhibit liver production or release of glucose in order to lower blood glucose in Type II diabetes (NIDDM) or to improve the profile of hyperinsulinemia and dyslipidemia in insulin resistance syndrome without diabetes. The mechanisms of action of pharmaceutical drugs and twelve herbal medicines that lower blood glucose are reviewed. Panax ginseng, Gymnema sylvestre, Brickellia spp Vaccinium spp and Syzygium jambolana appear to lower blood glucose through undesirable mechanisms. Panax quinquefolius, Foeniculum vulgare, Grifola spp Momordica charantia, Ocimum spp and Cinnamomum cassia may act at least in part by increasing insulin sensitivity in the tissues.

Introduction

Insulin resistance has been recognized as the root pathology underlying Type II diabetes (Non Insulin Dependent Diabetes Mellitus, or NIDDM) for decades, but only in recent years has insulin resistance syndrome (IRS) without loss of control of blood glucose been recognized as a major disease pathology in its own right. When the cells become resistant to the effects of insulin, the pancreas secretes larger amounts, and for a longer period of time. Figure 1 (page three) shows typical insulin curves after a glucose challenge in a variety of patients. The top curve is for an individual who may have perfectly normal fasting and post-meal blood glucose levels. The second curve shows a similar individual whose IRS has progressed to loss of glycemic control. This patient still has hyperinsulinemia, with the added pathological burden of elevated blood glucose. Both patients have elevated and prolonged insulin curves compared to the third curve, for an individual with both normal insulin and blood glucose levels. Research from the last decade

Continued on page three

Asthma case study

Chanchal Cabrera, M.N.I.M.H., M.Sc.

Abstract: A 36 year-old patient with asthma of 6 years duration was successfully weaned from prescriptions of Ventolin and Becloforte using a combination of herbal therapeutics and withdrawal of milk products from the diet. The patient was medication-free after several months and remained so for a year of follow-up.

Patient

Female. Aged 36. Presented with asthma which began 6 years ago during her 2nd pregnancy. She had a long history of repeated sinus infections and bronchitis and had received many antibiotics and steroids over the years. She was presently using Ventolin 5 - 10 puffs a day and Becloforte 2 puffs twice day. She felt a chest tightness all the time and all exertion was difficult. She had a lot of family stress and had a history of depression. There was a family history of eczema and allergies. On examination her chest expansion was greatly reduced and there was a slight expiratory wheeze. She had once been hospitalized for status asthmaticus and was afraid of this happening again.

Recommendations

She was advised to quit eating all dairy products and any foods with dairy in them and to avoid all sweetened and sugary foods and all processed or packaged foods. She was given information on avoiding dairy, reducing exposure to dust and airborne allergens, and on breathing exercises.

Supplements

Vitamin C to bowel tolerance
Evening Primrose oil 3 g./day
N-acetyl-cysteine 1500 mg./day
Quercitin 1 g./day

Continued on page fourteen

Insulin resistance *from page one*

has demonstrated that such elevated insulin, with accompanying metabolic factors, is a root cause of many of the diseases of modern civilization even in the absence of overt diabetes.

Pathophysiology of hyperinsulinemia

In normal metabolism, as insulin rises after a meal, or in response to stress-related hyperglycemia during periods of stress, it triggers release of insulin-like growth factor (IGF-1) from the liver. This in turn suppresses growth hormone, and IGF-1 effects combine with those of insulin for a net anabolic effect on cells. The combination of elevated or prolonged insulin secretion in IRS, the accompanying excessive and prolonged secretion of IGF-1, with extended suppression of growth hormone can have powerful pathological effects on the tissues. Some chronic conditions related to insulin resistance and compensatory hyperinsulinemia, but with normal glycemic control, are listed below.

Obesity

Hypertension

Atherosclerosis

Elevated triglycerides with depressed HDL cholesterol

Increasing the tendency of the blood to clot

Heart attacks and strokes

Polycystic ovarian syndrome

Elevated IGF-1 accompanying insulin resistance may also be related to the pathology of cancers of the breast, prostate, and colon.

All the above may occur in an individual with normal control of blood sugar. In NIDDM, elevated blood glucose may add to the damage, but simple reduction of blood glucose will not remedy the problem. Both insulin and blood glucose must be lowered, preferably by correcting the original resistance of the cells to the effects of insulin. Glycemic control in diabetes attained through stimulation of insulin in a patient who is already hyperinsulinemic may worsen the prognosis of the disease or threaten serious hypoglycemic episodes. Agents that suppress liver production or release of glucose have historically led to liver disease in some patients, and at least three leading drugs in this category had to be pulled from the market due to hepatotoxicity.

The patient above with the highest insulin curve has normal blood glucose. In the patient with NIDDM, therapies that normalize blood glucose by elevating insulin worsen the underlying pathology of diabetes. (Bagdade et al.)

Natural methods to increase insulin sensitivity

Drug or herbal treatments to correct insulin resistance and hyperinsulinemia are irrational as permanent cures, because the syndrome is easily treated with lifestyle and dietary modification. A combination of reduction of carbohydrates in the diet, regular moderate activity including cross training to tone multiple muscle systems, and supplementation with nutritional factors known to increase insulin rapidly improve measures of insulin resistance, including glycemic control in NIDDM. Some nutritional factors that may increase insulin sensitivity are listed below.

Chromium picolinate 200-400 mcg twice a day

Cod Liver Oil for a dose of docosahexaenoic acid (DHA) equivalent to 2-4 grams/day

B-Complex, in the form of a multivitamin

Magnesium 600-1000 mg/day

Zinc 30-60 mg/day

Drug actions

Drugs that lower blood glucose may do so in the following ways:

Reduce glucose absorption from the intestine. This must be viewed as a poor substitute for reducing carbohydrates in the diet.

Inhibit liver enzymes that release glucose from the liver. This may be either through suppression of the release of glucose from glycogen, or of manufacture of glucose through gluconeogenesis. Drugs of this class have potential liver toxicity as a side effect. This action does not address the underlying pathology of insulin resistance, but creates a disease condition in the liver to balance the disease condition in the cells.

Stimulate insulin production or release from the pancreas. This approach may lower blood glucose, but actually worsens the underlying pathology. The accompanying graph shows that insulin is already pathologically elevated in NIDDM. The hyperinsulinemia itself is pathogenic and produces many of the destructive symptoms of NIDDM. Stimulating it further may also hasten the ultimate exhaustion of the pancreas. Drugs which powerfully elevate insulin also increase the risk of fatal hypoglycemic events.

Increase insulin sensitivity. If a drug or herb is used for IRS or NIDDM, this should be the preferred effect. This should not, however, be expected to take the place of normal nutritional components or normal activity that decreases insulin resistance.

Pharmaceutical drugs

The quest for oral hypoglycemic agents was to prevent the inconvenience of insulin injections. For most of the later twentieth century the focus was on lowering blood glucose, without particular regard to the mechanism. With a growing understanding of insulin resistance as the underlying mechanism of a syndrome that is pathological in itself whether or not it includes loss of glycemic control, the focus in the recent decade has been to identify drugs that increase insulin sensitivity.

Guanidine derivatives

First oral hypoglycemic drugs were guanidine derivatives. These may have a combination effect of suppression of sugar production in the liver and increase in insulin sensitivity in the cells. The first drugs in the class, buformine and phenformine, were discontinued due to adverse effects on the liver. The current drug in this category is metformin (Glucophage).

Sulfonureas

The next generation of oral hypoglycemics are the sulfonureas. These increase pancreatic insulin, and symptomatically reduce blood sugar. These drugs can cause life-threatening hypoglycemia. They also elevate triglycerides, a normal consequence of hyperinsulinemia. They hasten the degeneration of the pancreatic insulin-producing cells in NIDDM. They are contraindicated in pre-diabetic insulin resistance due to the tendency to cause hypoglycemia.

Thiazolidinediones

Troglitazone (Rezulin) was widely heralded in the mid-nineties as an insulin-sensitizing agent, but eventually was withdrawn from the market after nearly 100 cases of acute Liver Failure developed in people taking it (Graham et al.).

Metformin

Metformin (Glucophage) is a guanidine derivative that inhibits liver production of glucose via gluconeogenesis, and also increases the insulin sensitivity of the peripheral tissues. It does not cause hyperinsulinemia or hypoglycemia and may be the most appropriate drug therapy where a drug is required to control blood glucose. It is widely promoted as an insulin sensitizer, but it also acts on the liver, has multiple side effects, and may cause death in individuals with impaired kidney function, a common condition in diabetes.

A study of representatives of the guanidine derivative and sulfonureas in 1971 found that although they improved control of blood glucose, they significantly increased deaths from cardiovascular causes relative to giving insulin alone (Cornfield; Schor; University Group Diabetes Program.).

Herbs

Many herbs have been used to treat diabetes or its symptoms in areas of the world and in eras when diabetes was present. Diabetes apparently first appeared in ancient Egypt, followed later in the grain farming valleys of India, China, and Mesopotamia. The first recorded case in European history appears to be in the 1600s in Holland. Some of these herbs reduce blood glucose, and others treat the symptoms of diabetes without lowering blood glucose. Considering current understanding of insulin resistance, the same discrimination should be used when selecting herbs that lower blood glucose as when selecting drugs. Herbs that

lower blood glucose by increasing insulin sensitivity may also be useful in IRS without diabetes. Many historical and currently popular herbs for lowering blood sugar do so through worsening the underlying state of hyperinsulinemia.

Panax

Various studies of both Asian and American ginseng have shown improvement in aspects of glycemic control in animals and humans. The mechanisms are not clear, but at least some constituents of *Panax ginseng* promote insulin production and release, and improvements in control of blood of glucose may be at the expense of stimulating hyperinsulinemia (Waki et al.; Kimura et al.).

A low dose of *P. ginseng* (100-200 mg) extract was given to 36 diabetic subjects for 8 weeks. The test groups showed improvements in fasting blood glucose and increases in daily activity and exercise. No mechanism was suggested for the activity, but there was no change in serum lipids in the ginseng group. If the action were through improving insulin sensitivity, a drop in serum triglycerides would be expected, so it appears that ginseng acts through methods other than the optimal increase of insulin sensitivity in the peripheral tissues (Sotaniemi et al.).

Oral administration of an oral preparation of unprocessed roots of *P. ginseng* in mice reduced blood glucose. The mechanisms suggested were through inhibition of intestinal glucose absorption, and suppressing liver production of glucose, the latter being an undesirable mechanism for long term use (Chung et al.).

A preparation of *P. ginseng* rootlets lowered blood glucose in animals, and the suggested mechanisms were inhibition of intestinal absorption of glucose, and an increase in insulin sensitivity. *P. ginseng* rootlets are considered a separate medicine from the root in Chinese herbalism, with rootlet actions resembling the yin tonic American ginseng (*Panax quinquefolius*) rather than the more stimulating chi and yang tonifying effects of *P. ginseng* (Chung et al.).

One gram doses of American ginseng reduced the area under the blood glucose curve in human volunteers without diabetes. The effect was statistically significant, but was slight, from 12-19% of the area under the curve. The ginseng had to be given at least 40 minutes before the glucose challenge to be effective, and a 3 gram dose provided no improved results over a 1 gram dose (Vuksan et al. 2001).

The same researchers found a clinically significant fall in the peak glucose level at the 2 hour point in patient with NIDDM, with administration of 3 gram of *P. quinquefolius* – blood glucose at the 2 hour point was 60% lower in the panax group than the placebo group (Vuksan et al. 2000b).

In an earlier trial, in both insulin-resistant diabetics and non-diabetic overweight individuals, the researchers found that a 3 gram dose given 40 minutes before a glucose challenge reduced the area under the post-meal glucose curve by just less than 20% (Vuksan et al. 2000a).

No mechanism has been suggested for the activity of American ginseng. The demonstrated insulin secretagogue effects constituents of *P.ginseng* roots suggest that it should be avoided in the treatment of insulin resistance. Insulin-sensitizing *P. ginseng* rootlets are generally not available in the U.S. herb market, but their traditional use as an analogue to *P. quinquefolius* at least suggests that *P. quinquefolius* should be used

Bastyr ad

in preference to *P. ginseng* for insulin resistance syndromes.

Gymnema

Gymnema sylvestre is a traditional Ayurvedic and Arabic medicine used for diabetes since antiquity in India (Nadkarni) and Egypt (El Gammel). Clinical trials have shown that it can lower blood sugar in both Insulin Dependent Diabetes (IDDM) and NIDDM (Shanmugasundaram et al.; Baskaran et al.). Typical conventional dose: 10 to 15 ml per day of a 1:1 extract or Gymnema tablets 4000 mg 3 daily. Research into the mechanism of action in animals suggests that the plant delays the absorption of glucose from the intestine (Yoshikawa et al.; Shimizu et al.). Consistent results in petrie dish, animal and human trials demonstrate that it increases insulin secretion (Persaud et al.), raises serum insulin (Shanmugasundaram et al.; Baskaran et al.), and does not act through increasing insulin sensitivity in the tissues (Tominaga et al.). Older research show that the plant can induce hypoglycemia in normal animals (Nadkarni). Thus gymnema has qualities in common with the sulfonureas drugs. It lowers blood sugar by worsening the underlying disease process of hyperinsulinemia.

Vaccinium

The leaves of *Vaccinium myrtillus* and related species (blueberry, bilberry) have been used to treat diabetes and its complications since antiquity in Egypt (El Gammel). They are also used in Central European folk medicine (Weiss; Petlevski). The use for diabetes in the U.S. appears to have been first recorded in the 1930s, when researchers found it used Swiss-American folk medicine, and verified a blood sugar lowering effect; by the 1960s it had entered into common usage as a treatment for elevated blood sugar in the U.S. via the health foods industry (Crellin and Philpott). In both traditional and contemporary literature, distinction has to be made between use of the leaf and the berry. Vaccinium berry and its flavonoid extracts are used to treat the vascular damage of diabetes without altering insulinemic or glycemic status (Savickiene et al.).

No studies appear to describe a mechanism for its action, but Weiss reports research showing that long-term use can cause severe hydroquinone poisoning and states that it is not appropriate for regular use. Weiss does not state the form or dose that may cause poisoning. Appalachian herbalist Tommie Bass states that the dose he recommended was several hands full

of the leaf in a gallon of water to make tea and makes no observation of toxicity (Crellin and Philpott).

Oplopanax

Oplopanax horridum (devil's club) has a reputation among contemporary herbalists as a plant that can lower blood sugar in diabetes. The first report of this historically was based on an anecdote in British Columbia where a native coming to a hospital for surgery suddenly developed the symptoms of diabetes. The physician enquired and it seems that the man had been taking devil's club. The doctor postulated that the devils club was effectively treating the diabetes, and when the plant was withdrawn in the hospital, the full disease emerged. The case was written up and published in a Canadian medical journal. The problem with this particular anecdote is that borderline NIDDM often appears only under conditions of stress, and then disappears when the stress is removed. An aboriginal man admitted to a hospital for anesthesia and surgery could have been experiencing severe stress, and this could have caused the emergence of the symptoms of NIDDM rather than withdrawal of the oplopanax. Subsequent clinical trials have found no effect of oplopanax on either blood sugar or insulin (Justice; Thommassen et al.).

A review of the older ethnobotanical literature on oplopanax turned up only a single reference to its use in diabetes: one native informant reported that she remembered a white woman taking the plant for diabetes (Thie). Subsequently, it appears that in recent decades use of the plant among natives for diabetes has become common (Thommassen et al.), and it appears from the published studies that the plant benefits patients subjectively in some way unrelated to glycemic control (Justice; Thommassen et al.).

Because oplopanax has no demonstrated action of blood glucose or insulin there is no rationale for using it to treat underlying insulin resistance syndrome.

Brickellia

Brickellia grandiflora (progidiosa) and related species are used in popular medicine in Mexico and the Southwest U.S. to control elevated blood sugar in diabetes (Moore). Research in Mexico suggests that the mechanism is through stimulation of insulin from the pancreas (Peres-Gutierrez). Extracts were given to diabetic and normal mice. As expected, the extract lowered blood sugar in the diabetic mice, but also lowered blood sugar by 40% in the normal mice. An agent acting to normalize insulin resistance would not

be expected to do this, while the result is typical of an agent that either stimulates insulin release or decreases liver release of glucose.

Syzygium

Syzygium jambolana, fructus (*S. cumini*, *Eugenia jambolana* Lam, *Eugenia cumini* Druce, *Myrtus cumini* Linn), common name Jambul, has been used in traditional Egyptian (El Gammel), Unani (Said), and Ayurvedic medicine (Kar et al.; Grover et al. 2001; Rathi) to treat diabetes. The activity has been confirmed in some animal trials (Kar et al.; Grover et al., 2001; Rao and Rao), but not in others. The plant part used may be important. The leaf decoction was found ineffective in diabetic animals (Pepato et al.) and in non-diabetic human volunteers (Teixeira et al. 2000). A tea of the seeds was ineffective in animals (Teixeira et al. 1997). Powders or decoctions of the dried seeds were effective in animals in another trial (Grover et al. 2000). The fruit pulp was effective in another animal trial (Achrekar et al.). Research into the mechanisms of action of *Syzygium* are contradictory, with some animal trials showing an inhibition of the liver enzymes that release blood glucose (Grover et al. 2000) or stimulating insulin secretion (Achrekar et al.), and one trial showed a lowered insulin response to a fructose challenge (Vikrant et al. 2001).

Herbs that may enhance insulin sensitivity

Trigonella

In a trial of a fenugreek extract (*Trigonella foenum-gracium*) in twenty-five patients with Type II diabetes, a trial group of thirteen patients received one gram a day of an evaporated hydro-alcoholic extract of fenugreek seeds for two months. At the end of the period, blood sugar responses to a meal were significantly lower in the fenugreek group. Insulin secretion was also lower, as were serum triglycerides (TG). HDL cholesterol was improved, and a standard measure of insulin sensitivity showed increased insulin sensitivity (Gupta et al.).

Dose of powder has varied in different trials from 18 grams (Sowmya and Rajyalakshmi) to 100 grams (Sharma 1990) with several trials giving 25 grams (Sharma, Sarkar et al.). Long term compliance can therefore be difficult. In some clinical trial in India, the powder was mixed into chipati flour for easier compliance. In several trials, an evaporated hydro-alcoholic

extract was effective (Gupta et al.), as was an aqueous extract (Abdel-Barry).

Cinnamomum

Chinese diabetes researchers investigating traditional formulas for diabetes discovered that a formula based on cinnamon and prepared aconite (*quei fu di huang wan*) lowered blood glucose in test animals. The formula suppressed the production of glucose in the liver in a manner similar to that of metformin. They then separately tested the cinnamon and the aconite. Both herbs lowered glucose, the aconite by suppressing gluconeogenesis, and the cinnamon by some other unidentified mechanism (Cheng et al.).

Studies at the U.S. Department of Agriculture suggest that cinnamon acts in some manner to potentiate the action of insulin on the target cell (Berrio et al.; Broadhurst et al.; Khan et al.). Other researchers found that cinnamon components help to activate specific cellular responses to insulin (Imparl-Radosevish et al.; Jarvill-Taylor et al.) thus increasing insulin sensitivity.

Grifola

Maitake mushroom (*Grifola spp.*) has been shown in animals to lower both blood glucose and insulin (Talpur; Manohar; Horio; Kubo). It improves many aspects of hyperinsulinemia, including blood sugar, HBA1C (a measure of long term glycemic control) (Talpur et al.), insulin levels, and triglycerides (Talpur et al.; Kubo et al.). Two case reports have appeared in the literature suggesting that grifola polysaccharides in moderate doses may completely replace conventional antidiabetic drugs (Konno). The animal trials indicate that either the powder of maitake or a water extract may be effective. The mechanism seems only to involve an increase in insulin sensitivity.

Momordica charantia

Momordica charantia (bitter melon) has been in continuous use since antiquity to treat diabetes in Ayurvedic, Unani, and Egyptian medicine (Grover et al. 2002; El Gammel). Many trials have verified its use to lower blood glucose in experimental animals (Kar; Miora; Vikrant), and the property has been confirmed in humans, where the plant lowered blood glucose in 86% of diabetic patients tested (Ahmad). It has also been shown to reduce diabetic pathological endpoints in animals, such as cataract (Rathi) and kidney damage (Grover et al., 2001). It appears to act by increasing insulin sensitivity in the peripheral tissues (Miura et al.).

Momordica prevents both hyperinsulinemia and hypertriglyceridemia (Vikrant et al.).

Ocimum

Various species of basil (*Ocimum* species) have been used in traditional medicine to treat diabetes, including *O. sanctum* (Kar et al.; Grover et al.), *O. canum* (Nyarko et al.), *O. gratissimum* (Aguiyi et al.) and *O. ablum* (Agrawal et al.). A number of trials have demonstrated a hypoglycemic effect in animals (Kar et al.; Nyarko et al.; Vats et al.; Aguiyi et al.; Chattopadhyay; Rai et al.). An anti-hyperglycemic effect has been confirmed on both fasting and post-prandial blood glucose in a human clinical trial (Agrawal et al.). Studies in animals (Chattopadhyay et al.) and humans (Viseshakul et al.) suggest that its action is through increasing insulin sensitivity in the cells.

Abdel-Barry JA, Abdel-Hassan IA, Jawad AM, al-Hakim MH. Hypoglycaemic effect of aqueous extract of the leaves of *Trigonella foenum-graecum* in healthy volunteers. **East Mediterr Health J** 2000 Jan;6(1):83-8

Achrekar S, Kaklij GS, Pote MS, Kelkar SM. Hypoglycemic activity of *Eugenia jambolana* and *Ficus bengalensis*: mechanism of action. **In Vivo** 1991 Mar-Apr;5(2):143-7

Agrawal P, Rai V, Singh RB. Randomized placebo-controlled, single blind trial of holy basil leaves in patients with noninsulin-dependent diabetes mellitus. **Int J Clin Pharmacol Ther** 1996 Sep;34(9):406-9

Aguiyi JC, Obi CI, Gang SS, Igweh AC. Hypoglycaemic activity of *Ocimum gratissimum* in rats. **Fitoterapia** 2000 Aug;71(4):444-6

Ahmad N, Hassan MR, Halder H, Bennoor KS. Effect of *Momordica charantia* (Karolla) extracts on fasting and post-prandial serum glucose levels in NIDDM patients. **Bangladesh Med Res Counc Bull** 1999 Apr;25(1):11-3

Bagdade JD, Bierman EL, Porter D. The significance of basal insulin levels in evaluation of the insulin response in diabetic and non-diabetic subjects. 1967. **J Clin Invest**(NY) 46(10), 1553.

Baskaran K, Kizar Ahamath B, Radha Shanmugasundaram K, Shanmugasundaram ER. Antidiabetic effect of a leaf extract from *Gymnema sylvestre* in non-insulin-dependent diabetes mellitus patients. **J Ethnopharmacol** 1990 Oct;30(3):295-300

Berio LF, Polansky MM, Anderson RA. Insulin activity: stimulatory effects of cinnamon and brewer's yeast as influenced by albumin. **Horm Res** 1992;37(6):225-9

Broadhurst CL, Polansky MM, Anderson RA. Insulin-like biological activity of culinary and medicinal plant aqueous extracts in vitro. **J Agric Food Chem** 2000 Mar;48(3):849-52

Chattopadhyay RR. Hypoglycemic effect of *Ocimum sanctum* leaf extract in normal and streptozotocin diabetic rats. **Indian J Exp Biol** 1993 Nov;31(11):891-3

Cheng JT, Liu IM, Chi TC, Su HC, Chang CG. Metformin-like effects of Quei Fu Di Huang Wan, a Chi-

nese herbal mixture, on streptozotocin-induced diabetic rat. **Horm Metab Res** 2001 Dec;33(12):727-32

Chung SH, Choi CG, Park SH. Comparisons between white ginseng radix and rootlet for antidiabetic activity and mechanism in KKAY mice. **Arch Pharm Res** 2001 Jun;24(3):214-8
Cornfield J. The University Group Diabetes Program. A further statistical analysis of the mortality findings. **JAMA**. 1971 Sep 20 ;21 7(12):1676-87.

Crellin JK and Philpott J. **A Reference Guide to Medicinal Plants: Herbal Medicine Past and Present**. Durham, North Carolina: Duke University Press, 1989

El Gammel SY Antidiabetic herbs in history, in: Adly A. [Ed] **The History of Medicinal Plants: Proceedings of the Second International Congress Organized by the Arab Society for the History of Pharmacy in co-operation with the World Union of Societies of the History of Pharmacy**. Alexandria Egypt, December 15th to 19th 1980. Karachi, Hamdard Foundation Press, 1980

Graham DJ, Drinkard CR, Shatin D. Incidence of idiopathic acute liver failure and hospitalized liver injury in patients treated with troglitazone. **Am J Gastroenterol** 2003 Jan;98(1):175-9

Grover JK, Yadav S, Vats V. Medicinal plants of India with anti-diabetic potential. **J Ethnopharmacol** 2002 Jun;81(1):81-100

Grover JK, Vats V, Rathi SS, Dawar R. Traditional Indian anti-diabetic plants attenuate progression of renal damage in streptozotocin induced diabetic mice. **J Ethnopharmacol** 2001 Aug;76(3):233-8

garden

- Grover JK, Vats V, Rathi SS. Anti-hyperglycemic effect of *Eugenia jambolana* and *Tinospora cordifolia* in experimental diabetes and their effects on key metabolic enzymes involved in carbohydrate metabolism. **J Ethnopharmacol** 2000 Dec;73(3):461-70
- Gupta A, Gupta R, Lal B. Effect of *Trigonella foenum-graecum* (fenugreek) seeds on glycaemic control and insulin resistance in type 2 diabetes mellitus: a double blind placebo controlled study. **J Assoc Physicians India** 2001 Nov;49:1057-61
- Said HM [Editor] **Hamdard Pharmacopoeia of Eastern Medicine**. Delhi; Sri Satguru Publications, 1970.
- Horio H, Ohtsuru M. Maitake (*Grifola frondosa*) improve glucose tolerance of experimental diabetic rats. **J Nutr Sci Vitaminol** (Tokyo) 2001 Feb;47(1):57-63
- Imparl-Radosevich J, Deas S, Polansky MM, Baedke DA, Ingebritsen TS, Anderson RA, Graves DJ. Regulation of PTP-1 and insulin receptor kinase by fractions from cinnamon: implications for cinnamon regulation of insulin signalling. **Horm Res** 1998 Sep;50(3):177-82
- Jarvill-Taylor KJ, Anderson RA, Graves DJ. A hydroxychalcone derived from cinnamon functions as a mimetic for insulin in 3T3-L1 adipocytes. **J Am Coll Nutr** 2001 Aug;20(4):327-36
- Justice, J.W. Use of Devil's Club in Southeast Alaska. **Alaska Medicine** 8(2) 36-39
- Kar A, Choudhary BK, Bandyopadhyay NG. Comparative evaluation of hypoglycaemic activity of some Indian medicinal plants in alloxan diabetic rats. **J Ethnopharmacol** 2003 Jan;84(1):105-8
- Khan A, Bryden NA, Polansky MM, Anderson RA. Insulin potentiating factor and chromium content of selected foods and spices. **Biol Trace Elem Res** 1990 Mar;24(3):183-8
- Kimura M, Waki I, Chujo T, Kikuchi T, Effects of hypoglycemic components in ginseng radix on blood insulin level in alloxan diabetic mice and on insulin release from perfused rat pancreas. **J Pharmacobiodyn** 1981 Jun;4(6):410-7
- Konno S, Tortorelis DG, Fullerton SA, Samadi AA, Hettiarachchi J, Tazaki H. A possible hypoglycaemic effect of maitake mushroom on Type 2 diabetic patients. **Diabet Med** 2001 Dec;18(12):1010
- Kubo K, Aoki H, Nanba H. Anti-diabetic activity present in the fruit body of *Grifola frondosa* (Maitake). **Biol Pharm Bull** 1994 Aug;17(8):1106-10
- Manohar V, Talpur NA, Echard BW, Lieberman S, Preuss HG. Effects of a water-soluble extract of maitake mushroom on circulating glucose/insulin concentrations in KK mice. **Diabetes Obes Metab** 2002 Jan;4(1):43-8
- Miura T, Itoh C, Iwamoto N, Kato M, Kawai M, Park SR, Suzuki I. Hypoglycemic activity of the fruit of the *Momordica charantia* in type 2 diabetic mice. **J Nutr Sci Vitaminol** (Tokyo) 2001 Oct;47(5):340-4
- Moore, M. **Medicinal Plants of the Desert and Canyon West**. Santa Fe: Museum of New Mexico Press, 1989
- Nadkarni KM **Indian Materia Medica**. Bombay: Popular Prakashan
- Perez-Gutierrez RM, Perez-Gonzalez C, Zavala-Sanchez MA, Perez-Gutierrez S. Hypoglycemic activity of *Bouvardia terniflora*, *Brickellia veronicaefolia*, and *Parmentiera edulis*. **Salud Publica Mex** 1998 Jul-Aug;40(4):354-8
- Pepato MT, Folgado VB, Kettelhut IC, Brunetti IL. Lack of antidiabetic effect of a *Eugenia jambolana* leaf decoction on rat streptozotocin diabetes. **Braz J Med Biol Res** 2001 Mar;34(3):389-95
- Persaud SJ, Al-Majed H, Raman A, Jones PM. *Gymnema sylvestre* stimulates insulin release in vitro by increased membrane permeability. **J Endocrinol** 1999 Nov;163(2):207-12
- Petlevski R, Hadzija M, Slijepcevic M, Juretic D. Effect of 'antidiabetis' herbal preparation on serum glucose and fructosamine in NOD mice. **J Ethnopharmacol** 2001 May;75(2-3):181-4
- Rai V, Iyer U, Mani UV. Effect of *Tulasi* (*Ocimum sanctum*) leaf powder supplementation on blood sugar levels, serum lipids and tissue lipids in diabetic rats. **Plant Foods Hum Nutr** 1997;50(1):9-16
- Rao BK, Rao CH. Hypoglycemic and antihyperglycemic activity of *Syzygium alternifolium* (Wt.) Walp. seed extracts in normal and diabetic rats. **Phytomedicine** 2001 Mar;8(2):88-93
- Rathi SS, Grover JK, Vikrant V, Biswas NR. Prevention of experimental diabetic cataract by Indian Ayurvedic plant extracts. **Phytother Res** 2002 Dec;16(8):774-7
- Savickiene N, Dagilyte A, Lukosius A, Zitkevicius V. Importance of biologically active components and plants in the prevention of complications of diabetes mellitus. **Medicina** (Kaunas) 2002;38(10):970-5
- Schor S. The University Group Diabetes Program. A statistician looks at the mortality results. **JAMA**. 1971 Sep 20;217(12):1671-5.
- Sowmya P, Rajyalakshmi P. Hypocholesterolemic effect of germinated fenugreek seeds in human subjects. **Plant Foods Hum Nutr** 1999;53(4):359-65
- Sharma RD, Raghuram TC, Rao NS. Effect of fenugreek seeds on blood glucose and serum lipids in type I diabetes. **Eur J Clin Nutr** 1990 Apr;44(4):301-6
- Sharma RD, Sarkar A, Hazra DK, Mishra B, et al. Use of fenugreek seed powder in the management of non-insulin dependent diabetes mellitus. **Nut. Res.** 16, 1331-1339
- Shanmugasundaram ER, Rajeswari G, Baskaran K, Rajesh Kumar BR, Radha Shanmugasundaram K, Kizar Ahmath B. Use of *Gymnema sylvestre* leaf extract in the control of blood glucose in insulin-dependent diabetes mellitus. **J Ethnopharmacol** 1990 Oct;30(3):281-94
- Shimizu K, Ozeki M, Tanaka K, Itoh K, Nakajyo S, Urakawa N, Atsuchi M. Suppression of glucose absorption by extracts from the leaves of *Gymnema inodorum*. **J Vet Med Sci** 1997 Sep;59(9):753-7
- Sotaniemi EA, Haapakoski E, Rautio A. Ginseng therapy in non-insulin-dependent diabetic patients. **Diabetes Care** 1995 Oct;18(10):1373-5
- Talpur NA, Echard BW, Fan AY, Jaffari O, Bagchi D, Preuss HG. Antihypertensive and metabolic effects of whole Maitake mushroom powder and its fractions in two rat strains. **Mol Cell Biochem** 2002 Aug;237(1-2):129-36
- Thie, K. **Medicinal Plants of the Pacific Northwest: A Digest of Anthropological Writings about Native**

References continued on back page

Asthma case from page one

Tincture Formula

<i>Ligusticum porteri</i> (1:4) (osha)	15
<i>Glycyrrhiza glabra</i> (1:3)(licorice)	15
<i>Angelica arch</i> (angelica) (1:4)	15
<i>Grindelia camporum</i> (1:3) (gumweed)	10
<i>Equisetum arvense</i> (1:3)(horsetail)	10
<i>Chamomilla recutita</i> (1:3) (chamomile)	10
<i>Urtica dioica</i> (1:3)(nettle)	10
<i>Euphorbia spp.</i> (1:4) (spurge)	10
<i>Lobelia inflata</i> (1:4) (lobelia)	5
<i>Eucalyptus</i> essential oil	4 drops
<i>Cinnamon</i> essential oil	4 drops
	100 mL

5 mL three times daily between meals

Tea formula: *Inula helenium* (elecampane), *Althea officinalis* (marshmallow leaf), *Glycyrrhiza glabra* (licorice), *Prunus serotina* (wild cherry bark), *Verbascum thapsus* (mullein), *Plantago lanceolata* (plantain), *Angelica archangelica* (angelica), *Ligusticum porteri* (osha), *Pimpinella anisum* (anise).

She was also given an essential oil blend to use for steam inhalations. Eucalyptus, Peppermint, Niouli, Camphor, Thyme.

Emergency tincture formula

Additionally I gave her an 'Emergency formula' to keep on hand in case she needed it.

<i>Viburnum opulus</i> (1:3)(cramp bark)	25
<i>Glycyrrhiza glabra</i> (1:4)(licorice)	20
<i>Ephedra sinica</i> (1:4)(ma huang)	20
<i>Datura stramonium</i> (1:5) (thorn apple)	15
<i>Valerian officinalis</i> (1:4)(valerian)	10
<i>Lobelia inflata</i> (1:4) (lobelia)	10
	100 mL

2.5 mL every 15 minutes as needed

Therapeutic rationale

The intention here was to strengthen the underlying lung function with equisetum, angelica and

glycyrrhiza while at the same time aiding in the removal of thick, sticky mucous with ligusticum, euphorbia and grindelia and reducing the allergic tendency with urtica and chamomilla. The lobelia and euphorbia were included to relax the smooth muscle of the lungs and reduce spasms and tightness of the airways.

The tea was prescribed as a gentle adjunctive support to the tincture. The essential oils are mucolytic and spasmolytic as well as warming and stimulating to lung function.

The nutritional supplements were prescribed to support the immune function and reduce the allergic reactions. For the same reasons she was asked to avoid all dairy products.

The emergency formula was given as an acute case remedy for symptomatic relief of occasional tightness and spasms in the respiratory system.

Outcome

She returned a month later and reported a dramatic improvement. She had used the Ventolin only twice in the past month and the Becloforte one puff twice a day. She had removed all dairy from her diet and significantly reduced the carbohydrates and sugars. She was walking 40 minutes a day without wheezing. Over the next few months she continued to improve and has not used any prescription medicines for over a year now.

Once we had the asthma under control, I reduced the tincture to once a day for 6 months and she now takes the tea 1 cup daily. She also took a tincture for 4 months to strengthen the immune system and now has not had a cold for 8 months.

<i>Allium sativum</i> (1:4) (garlic)	20
<i>Astragalus membr.</i> (1:3) (milk vetch)	20
<i>Tabebuia impetiginosa</i> (1:3) (taheebo)	20
<i>Calendula officinalis</i> (1:4) (marigold)	20
<i>Arctium lappa</i> (1:4)(burdock)	<u>20</u>
	100 mL

5 mL twice a day

This case was presented at the 1998 Clinical Case Studies Conference in Boulder, Colorado. The conference was presented by the Rocky Mountain Center for Botanical Studies and **Medical Herbalism**.

Traditional Medicine

Scullcap (*Scutellaria lateriflora*)

by Paul Bergner

More than a hundred species of scullcap (*Scutellaria spp.*) grow throughout the world. North American varieties were used by such native American tribes as the Penobscot, Iroquois, and Cherokee and by early European colonists for a variety of purposes. Several species grow in Britain, but do not appear in the classic British herbals of Culpepper and Gerard. *Scutellaria lateriflora* was used by all schools of North American medicine during the 1800s, has been in common use by herbalists in Great Britain from that time up to the present day. The plant is still used in rural folk medicine in the Appalachians, and is one of the most commonly used agents by contemporary medical herbalists in the U.S. and Britain. It was official in the *U.S. Pharmacopoeia* from 1860-1900.

Scullcap is useful for a general pattern of nervous exhaustion, from a variety of causes. When this condition is advanced, there is nervous depression with nervous agitation, restlessness, insomnia, and musculoskeletal tension and agitation. A more extreme pattern may include neuralgia, sciatica, tremors, and seizures. Scullcap has been used by physicians for such severe conditions as rabies and *delirium tremens*. Scullcap is not appropriate for acute or sub acute *inflammatory* conditions of the nervous system. Its particular pattern may underlie conditions of various organ systems, but if the pattern is remembered, then the application to specific conditions will be appropriate. The nervous exhaustion may result from acute or chronic disease, fevers, severe chronic cough, chronic poor digestion, acute or chronic reproductive complaints, overwork, teething in infants, caffeine abuse, episodes of mania or other mental illness, drug addiction or alcoholism, chronic sleep debt, overwhelming life events, chronic trauma and post traumatic stress disorder. The clinical presentation may include, besides fatigue, depression, nervous tension, and musculoskeletal agitation, a weak, tense pulse, palpitations or irregular heart beat, with tension headache or temporomandibular joint syndrome.

Scullcap has been classified as a nervine tonic and antispasmodic and used in that manner continuously since early in the 1800s. It appears to act on the central

nervous system to build and restore its tone, and paradoxically to allow relaxation and rest. This apparent contradiction – tonifying while relaxing – is the unique characteristic of scullcap, and the reason for its high esteem in North American medicine for more than 200 years. It is also mildly bitter and possesses the properties common to other bitter tonics with less tendency to overstimulate than agents such as gentian or hydrastis. It is sedative but does not leave a hangover the way valerian does in some doses for some patients. Due to volatile compounds, if the plant material is reasonably fresh and not overly heated during preparation, it possesses very mild diaphoretic and emmenagogue properties. Results may vary from patient to patient but generally scullcap does not possess strong hypnotic properties, and will not induce sleepiness during normal activities during the day. The most important exception is in the patient with chronic sleep debt, in whom scullcap may relax the system sufficiently that the normal urge to sleep is restored.

- For **general nervousness** combine with other nervine tonics such as tincture of milky oats.
- For **chronic stress** combine with herbs such as pulsatilla and vervain.

Continued on page seventeen

The Maker, in benevolence, has bestowed compound properties on by far the greater number of the agents placed at our disposal. Only a moderate number can be said to possess one property only. . . . In very many instances, one property is so predominant that the agent is valued chiefly for it, though possessing other properties . . . Taking in the vast number of our true remedies, considering the wide range of properties they occupy in this scale, and further remembering the diversity of organs on which they act and the diversity of times occupied in their action, and the mind will at once be astonished at the surprising sum of powers that have been provided for the

Historical combinations with Scullcap

Actions: bitter, nervine, tonic, antispasmodic, hypnotic

Nervine tonic

Scutellaria, Avena

For neurasthenia and neuroses Priest

Tonic nervine BHP

Add Turnera, Cola, and/or Humulus
for nervous disorders BHP

Add Verbena, with Cypripedium for
depressive states BHP

Scutellaria-Turnera

Tonic nervine Priest

Add verbena hastata, nervine tonic
With weak stomach Shook

Insomnia

Scutellaria-Passiflora

For insomnia Cook

Passiflora 15-40 drops; Scutellaria 2-12
drops for insomnia, exhaustion;
alcohol hangover Clymer

Add Humulus, Lactuca BHP

Scutellaria-Valeriana

Scutellaria 2; Valeriana 1 Bergner

Nervine antispasmodic

Scutellaria-Valeriana

Insomnia, antispasmodic Bergner

Add Pedicularis for tension
headache with TMJ Bergner

Add Viscum for hysteria BHP

Anxiety

Scutellaria-Turnera

For anxiety neurosis Priest

Add Verbena hastata, nervine tonic
with weak stomach, Shook

Neurological

Scutellaria-Stachys betonica

For nervous headache BHP

Add Cimicifuga for rheumatism or
chronic sciatica Priest

Scutellaria 1 oz.; Stachys elix 1.5 oz
Cimicifuga 1.5 oz; Helonias 1 oz; for
sciatica; 1 tsp per 2-3 hours with hot
sponge bath to affected parts Lyle

Scutellaria, Anemone, Cimicifuga

For convulsions Cook

For chorea, hysteria, agitation,
epeleptiform seizures Priest

Cough, fever, respiratory infection

Scutellaria-Asclepias tuberosa

Scutellaria 2-20 drops, Asclepias
15-40 drops per hour, for pneumonia,
acute pleurisy Clymer

Add Capsicum for influenza, colds, fever; chills
predominant Clymer

Add Lobelia for influenza, colds, fever; if relaxant
effect is needed Clymer

Scutellaria 2-15, Asclepias 20-45,
Marrubium 5-40 drops for deep seated
colds and cough with expectoration Clymer

Scutellaria 12-15 drops, Asclepias tub
20-40 drops; Eupatorium perf
15-30 drops for influenza, colds, fever,
pleurisy Clymer

Female, Pregnancy

Scutellaria, Caulophyllum

Restlessness during pregnancy Priest

Scutellaria, Caulophyllum 2.
Dioscorea 3; for convulsions, after-pains,
nervousness during parturition Lyle

Scutellaria 1, Caulophyllum 2,
Dioscorea 3; Lobelia (a little) for rigid
os uteri Lyle

Sources: *The Physiomedical Dispensatory*, Wm Cook, 1869; *Physiomedical Therapeutics, Materia Medica, and Pharmacy*, by T.J. Lyle, 1897; *Nature's Healing Agents*, RS Clymer, 1908; *Advanced Treatise in Herbology*, E. Shook; *Herbal Medication*, Priest and Priest, 1982; *British Herbal Pharmacopoeia*, 1983

The admixture of remedies should not be left to crude accident, but should be made a question of close scientific investigation; for the value of numerous articles can be greatly enhanced by giving them in suitable company.

Wm Cook 1869

- For **chronic insomnia**, combine with hypnotic-nervines and increase the dose approaching bedtime. Formula: scullcap, blue vervain, hops, and chamomile, with one half part of valerian.
- Take alone and with appropriate carminatives or demulcents for **nervous dyspepsia**.
- Combine with musculoskeletal remedies such as black cohosh, peony, kava, and/or pedicularis for **antispasmodic** effects.
- Combine with anodynes such as willow bark, California poppy, and wild lettuce, for exhaustion and insomnia accompanying **chronic pain**.
- Combine with relaxant expectorants such as wild cherry bark, pleurisy root, and lobelia for the exhaustion of **persistent acute cough**.
- Scullcap may help to relieve the restlessness and insomnia in **fever**. In the later stages of the fever, combine with relaxant diaphoretics such as asclepias, *Eupatorium perfoliatum*, or elder.
- For **fever in children**, combine with combination catnip and/or elder flower.
- For **convalescence from fevers** take alone or with stronger tonics.
- For **intermittent fevers**, combine with boneset (*Eupatorium perfoliatum*).
- For **teething in infants** combine with catnip.
- For **spasmodic uterine complaints** with general weakness and tendency to amenorrhea, combine with appropriate female system herbs.
- For **opium or alcohol withdrawal**, give larger doses of a strong infusion every two hours.
- For **tension headache** arising from chronic pain or coughing, use relaxants such as asclepias, pedicularis, kava, and/or peony.
- Scullcap has traditionally been used for **seizure disorder**, from a variety of causes, but traditional literature warns that it is fully effective or curative.
- The use of scullcap in **hydrophobia** was first commended by a Dr. Vandesveer in 1772, and

medical botanist Constantine Rafinesque reported on a large number of cured cases in 1830. The Eclectic John Uri Lloyd later documented cured cases and called for further formal research but the plant has not been formally studied for hydrophobia in either humans or animals.

Preparations

Traditional literature generally recommends a strong infusion (one ounce of good quality herbs per pint of water). William Cook (1868) warns that boiling destroys the medicinal properties of scullcap, and Scudder (1870) cautions that fresh or recently dried herbs should be used. As late as the 1920's The Eclectic Finley Ellingwood stated: "Its specific nerve sedative properties were those observed by the older writers who obtained this influence from a strong infusion which without doubt will yield results not obtained from small doses of the finer pharmaceutical preparations." The dose is about 4 ounces of the above infusion. The dose of the powder in the 1900 U.S.P. was 1 gram.

The accompanying table (page 16) gives some historical combinations in the Physiomedicalist tradition of herbalism.

medherb.com

- Online herbal bookstore with book reviews
- Annotated links
- Adverse Effects Forum
- Discussion forum
- Search site by keywords
- Electronic newsletter
- Online plant walk
- Online subscription renewal

Classifieds

Alchemy Botanicals is an herbal apothecary in Ashland, Oregon. Over 300 organic and wildcrafted bulk herbs, essential oils. Wide selection of herbal skincare products and medicine making supplies. Herbal formulations from Herbalist and Alchemist, Avena Botanicals, Wise woman Herbals, many others. Over 200 of the finest quality bulk tinctures and glycerites for you to create your own medicinal formulas. Practitioner discounts available. Call today for a free catalog. Shipping anywhere. Telephone 541-488-4418. Fax 541-488-4419. Email: herbs@alchemybotanicals.com. Alchemy Botanicals, 253 E Main Street, Ashland, OR 97520. Open 7 days a week.

Blue Ridge School of Herbal Medicine. Offers a 5-month, 250-hour Herbal Training Program focusing on holistic Chinese/Western diagnosis and the self-sufficient arts of plant identification and medicine making. Also offering a weekend exploration of Appalachian plants, "Herbal Safari" April 25-27, 2003. (828) 275-6221, coreypine@bigfoot.com, or P.O. Box 213, Asheville, NC 28802

Discover herbal medicine with Wise woman Ways. You will love Susun Weed's healing philosophy. Workshops near you, live-out apprenticeships, shamanic apprenticeships, correspondence courses, books, and videos. Information \$1. P.O. Box 64MH Woodstock, NY 12498

Healthcalls.net website for herbal education and protocols for today's most challenging health issues: porphyria, Lyme, detoxing, balancing Th1 and Th2 cytokines, antibiotic alternatives using essential oils, and training videos. Hart Brent, HealthCalls, 4287 Bayley Hazen Rd. West Danville, VT 05873 (802) 684-2570

Northeast School of Botanical Medicine. Comprehensive, practical training in clinical herbalism. Four programs offered. A six-month three days per week program; a seven month one weekend monthly course; Advanced Western Herbalism Intensive, and a traditional apprenticeship. Main instructor is 7song. Classes focus on constitutional diagnosis, plant identification, materia medica, first aid skills, student clinic, field trips, medicine preparation and formulation, anatomy and physiology, and community centered herbalism. P.O. Box 6626. Ithaca, NY 14851. 607-539-7172. Www.7song.com

Australasian College of Herbal Studies, USA Home Study and Residential Courses in Aromatherapy, Nutrition, Herbal Medicine, Homeopathy, Flower Essences, Homeobotanical Therapy, and Iridology. State Licensed. CE Credits available to RNs, Pharmacists, Veterinarians, and LMTs. Sallie Mae Student Loans. Veterans funding for on-campus programs. Free Program Catalog. 800-487-8839, achs@hrbed.com, www.herbed.com

The Herb Growing and Marketing Network The Herb Growing and Marketing Network. An information service for herb businesses and serious hobbyists. Includes The Business of Herbs, a 40 page trade journal; The Herbal Green Pages, annual resource guide with over 6000 listings; free classified advertising and lots of other benefits. Entire package \$95/yr. Sample journal \$6. HGMN, PO Box 245, Silver Spring, PA 17575-0245; 717-393-3295; herbworld@aol.com; <http://www.herbworld.com> and www.herbnet.com

HerbalGram. Highly acclaimed quarterly journal is now a benefit of membership in the American Botanical Council starting at \$50/year for individuals. Call 512-926-4900 or visit www.herbalgram.org

The Herb Quarterly. A beautiful magazine dedicated to all things herbal—gardening, medicinals, crafts, folklore, alternative uses of herbs, and more. Rates: \$24/1 yr, \$45/2 yrs, \$60/3 yrs. Box 548MH, Boiling Springs, PA 17007.

Classifieds are \$1 per word, \$35 minimum

Medical Herbalism

Senior Editor

Paul Bergner

Editor-in-Chief

Jill Hoppe, CCH

Associate Editors

Sharol Tilgner, ND
Steven Dentali, PhD

Contributing Editors

Chanchal Cabrera, MNIMH
Deborah Frances, ND
Jill Stansbury, ND

Editorial and Advertising Offices

P.O. Box 20512
Boulder, Colorado 80308

Phone

303-541-9552
303-415-9196 (fax)

Internet

bergner@medherb.com
<http://medherb.com>

Copyright 2003 North American Institute of Medical Herbalism, Inc

Adverse effects references from page nine

American Uses White Salmon, Washington: Longevity Herb Press, 1999

Thommassen HV, Wilson RA, McIlwain RG. Effect of devil's club on blood glucose levels in diabetes mellitus. **Canadian Family Physician** (1990) 36

Teixeira CC, Rava CA, Mallman da Silva P, Melchior R, Argenta R, Anselmi F, Almeida CR, Fuchs FD. Absence of antihyperglycemic effect of jambolan in experimental and clinical models. **J Ethnopharmacol** 2000 Jul; 71(1-2): 343-7

Teixeira CC, Pinto LP, Kessler FH, Knijnik L, Pinto CP, Gastaldo GJ, Fuchs FD. The effect of *Syzygium cumini* (L.) skeels on post-prandial blood glucose levels in non-diabetic rats and rats with streptozotocin-induced diabetes mellitus. **J Ethnopharmacol** 1997 May;56(3):209-13

Tominaga M, Kimura M, Sugiyama K, Abe T, Igarashi K, Igarashi M, Eguchi H, Sekikawa A, Ogawa A, Manaka H, et al. Effects of seishin-renshi-in and *Gymnema sylvestris* on insulin resistance in streptozotocin-induced diabetic rats. **Diabetes Res Clin Pract** 1995 Jul;29(1):11-7

University Group Diabetes Program Study pertaining to phenformin. **JAMA**. 1971 Aug 9;217(6):817.

Vats V, Grover JK, Rathi SS. Evaluation of anti-hyperglycemic and hypoglycemic effect of *Trigonella foenum-graecum* Linn, *Ocimum sanctum* Linn and *Pterocarpus marsupium* Linn in normal and alloxanized diabetic rats. **J Ethnopharmacol** 2002 Jan;79(1):95-100

Vikrant V, Grover JK, Tandon N, Rathi SS, Gupta N. Treatment with extracts of *Momordica charantia* and *Eugenia jambolana* prevents hyperglycemia and

hyperinsulinemia in fructose fed rats. **J Ethnopharmacol** 2001 Jul;76(2):139-43

Viseshakul D, Premvatana P, Chularojmontri V, Kewsiri D, Tinnarat P. Improved glucose tolerance induced by long term dietary supplementation with hairy basal seeds (*Ocimum canum sim*) in diabetics. **J Med Assoc Thai** 1985 Aug;68(8):408-11

Vuksan V, Sievenpiper JL, Wong J, Xu Z, et al. American ginseng (*Panax quinquefolius* L.) attenuates postprandial glycemia in a time-dependent but not dose-dependent manner in healthy individuals. **Am J Clin Nutr** 2001 Apr;73(4):753-8

Vuksan V, Sievenpiper JL, Koo VY, Francis T American ginseng (*Panax quinquefolius* L) reduces postprandial glycemia in nondiabetic subjects and subjects with type 2 diabetes mellitus. **Arch Intern Med** 2000(b) Apr 10;160(7):1009-13

Vuksan V, Stavro MP, Sievenpiper JL, Beljan-Zdravkovic U, et al. Similar postprandial glycemic reductions with escalation of dose and administration time of American ginseng in type 2 diabetes. **Diabetes Care** 2000 (b) Sep;23(9):1221-6

Waki I, Kyo H, Yasuda M, Kimura M. Effects of a hypoglycemic component of ginseng radix on insulin biosynthesis in normal and diabetic animals. **J Pharmacobiodyn** 1982 Aug;5(8):547-54

Weiss, RF. **Weiss's Herbal Medicine: Classic Edition**. Stuttgart:Theime, 2001

Yoshikawa M, Murakami T, Kadoya M, Li Y, Murakami N, Yamahara J, Matsuda H. Medicinal foodstuffs. IX. The inhibitors of glucose absorption from the leaves of *Gymnema sylvestris* R. BR. (Asclepiadaceae): structures of gymnemosides a and b. **Chem Pharm Bull** (Tokyo) 1997 Oct;45(10):1671-6

Subscription and Order form

(Fill in as many items as appropriate)

Name _____

Company _____

Street _____

City _____

State _____ Zip/Postal Code _____

Country _____

Telephone _____

Fax _____

Email _____

Visa/MC/Amex _____

Expiration Date ____ / ____

Call 303-541-9552

Medical Herbalism

P.O. Box 20512, Boulder, CO 80308

Http://medherb.com

U.S. Subscription

____ 1 year \$39.00 ____ 2 years \$70.00

Canada and Mexico

____ 1 year \$39.00 ____ 2 years \$70.00

North American Student

____ 1 year \$25.00 (photocopy of student i.d.)

Overseas

____ 1 year \$45.00 ____ 2 years \$80.00

Back Issues

(See pages 10-11)

____ Back issues at \$16.00 each = \$ ____ Total

Specify back issue numbers _____

Total order _____

(Cash, check, money order or charge in U.S.\$)